arXiv:1402.5296v1 [math.GR] 21 Feb 2014

LEFT TRANSITIVE AG-GROUPOIDS

M. RASHAD, I. AHMAD, AND M. SHAH

Abstract. An AG-groupoid is an algebraic structure that satisfies the left invertive law: $(ab)c = (cb)a$. We prove that the class of left transitive AG-groupoids (AG-groupoids satisfying the identity, $ab \cdot ac = bc$) coincides with the class of T^2-AG-groupoids. We also develop a simple procedure to test whether an arbitrary groupoid is left transitive AG-groupoid or not. Further we prove that, (i). Every left transitive AG-groupoid is transitively commutative AG-groupoid (ii) For left transitive AG-groupoid the properties of flexibility, right alternativity, AG*, right nuclear square, middle nuclear square and commutative semigroup are equivalent.

1. introduction

An AG-groupoid is an algebraic structure that satisfies the left invertive law, $(ab)c = (cb)a$[1]. This structure is a generalization of commutative semigroups. Every AG-groupoid satisfies the medial law, $(ab)(cd) = (ac)(bd)$. An AG-groupoid S with the left identity element e is called an AG-monoid. Every AG-monoid is paramedial, $(ab)(cd) = (db)(ca)$. Recently many researchers have been motivated to the field and a considerable work has been done on various aspects. Various classes have been discovered and enumerated up to order 6 [5, 6, 10, 11]. AG-groupoids have applications in the theory of flocks [1] and some in geometry[4].

A groupoid G satisfying the identity $ac \cdot bc = ab \ \forall a, b, c \in G$ is called right transitive groupoid [2], and G is called left transitive groupoid if it satisfying the identity $ab \cdot ac = bc$[13]. In Section 2 we prove that nonassociative right transitive AG-groupoids do not exist. In Section 3 we prove the coincidence of the left transitive AG-groupoids with the class of T^2-AG-groupoids. We prove that every left transitive AG-groupoid is transitively commutative AG-groupoid. We also investigate that on what conditions some left transitive AG-groupoids become commutative semigroups.

The following table contains various AG-groupoids with their defining identities that will be used in the rest of this article.

Key words and phrases. AG-groupoid, commutative semigroup, nuclear square AG-groupoid, unipotent AG-groupoid, T^2-AG-groupoid, alternative AG-groupoid.

AG-groupoid	defining identities
left nuclear square AG-groupoid	$a^2(bc) = (a^2b)c$
middle nuclear square AG-groupoid	$(ab^2)c = a(b^2c)$
right nuclear square AG-groupoid	$(ab)c^2 = a(bc^2)$
unipotent AG-groupoid	$a^2 = b^2$
T^1-AG-groupoid	$ab = cd \Rightarrow ba = dc$
T^2-AG-groupoid	$ab = cd \Rightarrow ac = bd$
T^3_l-AG-groupoid	$ab = ac \Rightarrow ba = ca$
T^3_r-AG-groupoid	$ba = ca \Rightarrow ab = ac$
T^3-AG-groupoid	both $T^3_l$and T^3_r
AG**-AG-groupoid	$a(bc) = b(ac)$
transitively commutative AG-groupoid	$ab = ba \,\&\, bc = cb$ $\Rightarrow ac = ca$
AG-monoid	an AG-groupoid with left identity
quasi-cancellative AG-groupoid	$x^2 = xy \;\&\; y^2 = yx \Rightarrow x = y$ $x^2 = yx \;\&\; y^2 = xy \Rightarrow x = y$
flexible AG-groupoid	$ab \cdot a = a \cdot ba$
right alternative AG-groupoid	$b \cdot aa = ba \cdot a$
AG*-groupoid	$(ab)c = b(ac)$

2. Right transitive AG-groupoids

We begin this section with the following definition of right transitive AG-groupoids and prove that nonassociative AG-groupoid does not exist.

Definition 1. *An AG-groupoid S is called right transitive AG-groupoid if $ab \cdot cb = ac \; \forall a, b, c \in S$.*

Theorem 1. *Every right transitive AG-groupoid is commutative semigroup.*

Proof. Let S be a right transitive AG-groupoid, and let $a, b \in S$. Then $ab = ab \cdot bb = (ab \cdot bb)(bb) = (bb \cdot bb)(ab) = bb \cdot ab = ba$. i.e. $ab = ba$. Thus S is commutative AG-groupoid and hence is associative. ∎

Due to their associative nature we avoid to discuss right transitive AG-groupoids in detail.

3. left transitive AG-test

Definition 2. *An AG-groupoid S is called left transitive AG-groupoid if,*

$$\forall a, b, c \in S, \; ab \cdot ac = bc. \tag{3.1}$$

We introduce a test to check an arbitarary AG-groupoid $(G, \cdot)$ for left transitive AG-groupoid. For this we define the following binary operation for some fixed $x \in G$ and $a, b \in G$.

$$a \star b = xa \cdot xb \tag{3.2}$$

(3.1) holds if,

$$a \star b = a \cdot b \tag{3.3}$$

To construct table of operation " $\star$ " for any fixed $x \in G$, we rewrite x-row of the " $\cdot$ " table as an index row of the new table and multiplying it by the elements of that row that is x-row of the " $\cdot$ " table from the left. If the tables for the operation " $\star$ " for each $x \in G$ coincides with the " $\cdot$ " table then 3.3 holds and hence the AG-groupoid is left transitive AG-groupoid. To illustrate the procedure, in the extended table for the following example, the tables on the right of the " $\cdot$ " table is obtained for the operation " $\star$ ".

Example 1. *Check the following AG-groupoids for left transitive AG-groupoid.*

$\cdot$	1	2	3
1	1	2	3
2	3	1	2
3	2	3	1

(i)

$\cdot$	1	2	3
1	1	1	1
2	1	1	3
3	1	2	1

(ii)

(i). We extend Table (i) in the way as described above to get the following:

$\cdot$	1	2	3	1	2	3	3	1	2	2	3	1
1	1	2	3	1	2	3	1	2	3	1	2	3
2	3	1	2	3	1	2	3	1	2	3	1	2
3	2	3	1	2	3	1	2	3	1	2	3	1

It is clear from the extended table that tables on the right coincide with the " $\cdot$ " table. Hence G is left transitive AG-groupoid.

(ii). To check Table (ii) for the desired property we extend the table as described.

·	1	2	3	1	1	1	1	1	3	1	2	1
1	1	1	1	1	1	1	1	1	1	1	1	1
2	1	1	3	1	1	1	1	1	1	1	1	1
3	1	2	1	1	1	1	1	1	1	1	1	1

The tables on the right do not coincide with the " $\cdot$ " table. Hence G is not left transitive AG-groupoid.

4. characterization of left transitive AG-groupoids

In this section we will discuss various relations of left transitive AG-groupoids with already known classes of AG-groupoids. We start with the following simple lemma which proves that the class of T^2-AG-groupoids (left transitive AG-groupoid) is contained in the class of unipotent AG-groupoids. This fact further assists to prove that every left transitive AG-groupoid is T^2-AG-groupoid and vice versa. For detailed study of T^2-AG-groupoids we suggest [8, 11].

Lemma 1. *Every T^2-AG-groupoid is unipotent AG-groupoid.*

Proof. Let S be a T^2-AG-groupoid and let b be any element of S. Then using the definition of T^2, we have $\forall a, b \in S$, $ab = ab \Rightarrow aa = bb \Rightarrow a^2 = b^2$. Hence S is unipotent AG-groupoid. ∎

Corollary 1. *Every unipotent AG-groupoid has a unique idempotent element.*

Using the above fact we are ready to prove that the class of T^2-AG-groupoids is the same as our class of left transitive AG-groupoids.

Theorem 2. *An AG-groupoid S is left transitive AG-groupoid if and only if it is T^2-AG-groupoid.*

Proof. $\Longrightarrow$ Let S be a left transitive AG-groupoid and $a, b, c, d \in S$. Assume that $ab = cd$. Then since S is left transitive we have $ac = ba \cdot bc = (ab \cdot aa)(bc)$ which implies by left invertive law that $ac = (bc \cdot aa)(ab)$. Now repeatedly using the medial and left invertive laws and the assumption $ab = cd$ we get, $ac = (bc \cdot c)(aa.d) = (cc \cdot b)(aa.d) = (cc \cdot aa)(bd) = (ca \cdot ca)(bd)$. Thus the left transitivity in S and the medial law further implies that $ac = aa \cdot bd = ab \cdot ad = bd \Rightarrow ac = bd$. Hence S is T^2-AG-groupoid.

Conversely, let S be a T^2-AG-groupoid and let $a, b, c \in S$. Then by Lemma 1 there exist c in S such that $a^2 = c^2 = c \; \forall \; a, c \in S$. Now by medial law $ab \cdot ac = aa.bc$ this implies by Lemma 1 and Corollary 1 that $ab \cdot ac = c \cdot bc$ $\Rightarrow$ $aa \cdot bc = c.bc$ (by medial law). This again by Corollary 1 implies that $bb \cdot bc = c.bc$ which by the left invertive law becomes $(bc \cdot b)b = c.bc$. This further implies by the use of the T^2-AG-groupoid and the left invertive law

alternatively that $(bc \cdot b)c = b \cdot bc \Rightarrow cb \cdot bc = b.bc \Rightarrow cb \cdot b = bc.bc \Rightarrow bb \cdot c = bc \cdot bc$. Thus by Lemma 1 and Corollary 1 we get $cc = bc \cdot bc$. Hence S is left transitive AG-groupoid. ∎

Thus left transitive AG-groupoids and the previous T^2-AG-groupoids coincide. Since we know the facts that every T^2-AG-groupoid is T^1-AG-groupoid, each T^1-AG-groupoid is T^3-AG-groupoid, T^1-AG-groupoid is paramedial and AG**-groupoid[8, 11, 12], thus we have the following corollary in terms of transitive AG-groupoids.

Corollary 2. *Every left transitive AG-groupoid is:*

(i) T^1-AG-groupoid
(ii) AG**-groupoid
(iii) T^3-AG-groupoid
(iv) Paramedial AG-groupoid
(v) Left nuclear square AG-groupoid
(vi) Unipotent AG-groupoid

Theorem 3. *Every left transitive AG-groupoid is transitively commutative AG-groupoid.*

Proof. Let S be a left transitive AG-groupoid and let $a, b, c \in S$. Assume that $ab = ba$, & $bc = cb$. Then since S is left transitive we have, $ac = ba \cdot bc$. This by use of left invertive law and assumption implies, $ac = (bc \cdot a)b = (cb \cdot a)b$ and once again using the left invertive law and the assumption we get $ac = (ab \cdot c)b = (ba \cdot c)b = bc \cdot ba$. But since S is left transitive we have $ac = ca$. Hence S is left transitively commutative AG-groupoid. ∎

The converse of the previous theorem is not true. We give a counterexample to show this.

Example 2. *Transitively commutative AG-groupoid of order* 4 *which is not a left transitive AG-groupoid.*

*	1	2	3	4
1	2	3	1	4
2	4	1	3	2
3	3	2	4	1
4	1	4	2	3

The following examples show that neither left transitive AG-groupoid nor AG-monoid is quasi-cancellative AG-groupoid. However, we can prove that the intersection of these two classes indeed becomes quasi-cancellative AG-groupoid.

Example 3. *(i) Left transitive AG-groupoids of order* 4 *which is not quasi-cancellative AG-groupoid. (ii) AG-monoid of order 4 which is not quasi-cancellative AG-groupoid.*

*	1	2	3	4
1	1	2	2	4
2	4	1	1	2
3	4	1	1	2
4	2	4	4	1

(i)

*	1	2	3	4
1	1	1	1	1
2	1	1	1	3
3	1	1	1	2
4	1	2	3	4

(ii)

Theorem 4. *Every left transitive AG-monoid is quasi-cancellative AG-groupoid.*

Proof. Let S be a left transitive AG-monoid and $a, b, c \in S$. For quasi-cancellative AG-groupoid, we have to prove that S is both left quasi-cancellative AG-groupoid and right quasi-cancellative AG-groupoid. For left quasi-cancellativity, let $a \cdot a = a \cdot b$ and $b \cdot b = b \cdot a$. Then $a = ea = be \cdot ba = be \cdot bb = eb = b \Rightarrow a = b$. Thus S is left quasi-cancellative AG-groupoid. Similarly it can be proved that S is right quasi-cancellative AG-groupoid. Hence S is quasi-cancellative and the theorem is proved. ∎

In the following we see that on what conditions a left transitive AG-groupoid becomes a semigroup.

Theorem 5. *For every left transitive AG-groupoid S the following are equivalent.*

(i) S is flexible AG-groupoid
(ii) S is right alternative AG-groupoid
(iii) S is AG*-groupoid
(iv) S is middle nuclear square AG-groupoid
(v) S is right nuclear square AG-groupoid
(vi) S is commutative semigroup

Proof. Let S be a left transitive AG-groupoid and $a, b, c \in S$.

$(i) \Rightarrow (ii)$. Assume that *(i)* holds. Now by left invertive law $ab \cdot b = bb \cdot a$. Since S is left transitive, this gives $ab \cdot b = (ab \cdot ab)a$. Which by left invertive law implies that$ab \cdot b = (a \cdot ab)(ab)$. The left transitivity and medial law further imply that $ab \cdot b = (a \cdot ab)(ba \cdot bb) = (a \cdot ba)(ab \cdot bb)$. We then have by assumption *get* that $ab \cdot b = (ab \cdot a)(ab \cdot bb)$. But since S is left transitive, we have $ab \cdot b = a \cdot bb$. Thus S is right alternative and hence *(ii)* holds.

$(ii) \Rightarrow (iii)$. Now assume that *(ii)* holds. Since by left inveritve law $ab \cdot c = cb \cdot a$. As S is left transitive, so by this and by left invertive law we have $ab \cdot c = (bc \cdot bb)a = (a \cdot bb)(bc)$. Thus by using assumption we have $ab \cdot c = (ab \cdot b)(bc)$. This further implies by the left transitivity in S and by medial law that $ab \cdot c = (ab \cdot b)(ab \cdot ac) = (ab \cdot ab)(b \cdot ac) = (bb)(b \cdot ac) = b \cdot ac \Rightarrow ab \cdot c = b \cdot ac$. Thus S is AG*-groupoid and hence *(iii)* holds.

$(iii) \Rightarrow (iv)$. Assume that *(iii)* holds i.e. S is AG*-groupoid, then $ab^2 \cdot c = b^2 \cdot ac$. Since S is left transitive this implies that $ab^2 \cdot c = b^2(ba \cdot bc)$ and by assumption again this implies that $ab^2 \cdot c = (ba \cdot bb)bc$. Again since S is left transitive and by assumption we have $ab^2 \cdot c = ab \cdot bc = (b \cdot ab)c = (ab \cdot b)c$. Thus by left invertive law and assumption we have $ab^2 \cdot c = (bb \cdot a)c = (b^2 \cdot a)c = a \cdot b^2c$. Hence $ab^2 \cdot c = a \cdot b^2c$ and thus S is middle nuclear square i.e. *(iv)* holds.

$(iv) \Rightarrow (v)$. Next assume that S is middle nuclear square AG-groupoid. Since S is left transitive we have $a \cdot bc^2 = a(ab \cdot ac^2)$, this implies by medial law that $a \cdot bc^2 = a(aa \cdot bc^2)$, which by assumption implies $a \cdot bc^2 = aa^2 \cdot bc^2$. This by the twice use of medial law gives $a \cdot bc^2 = ab \cdot a^2c^2 = ab(ac \cdot ac)$. But since S is left transitive we have $a \cdot bc^2 = ab \cdot c^2$. Hence S is right nuclear square and thus *(v)* holds.

$(v) \Rightarrow (vi)$. Assume that *(v)* holds. Since S is left transitive, we have $ab = aa \cdot ab = (ba \cdot ba)(a \cdot b) = (ba \cdot ba)(ba \cdot bb)$, which by assumption implies that $ab = ((ba \cdot ba)ba)bb$. This further implies by left invertive law that $ab = (bb \cdot ba)(ba \cdot ba)$. Thus by left transitive and medial properties in S we have $ab = ba(ba \cdot ba) = ba(bb \cdot aa)$, but since S is right nuclear square by assumption we have $ab = (ba \cdot bb)aa$. This by left transitivity implies that $ab = ab \cdot aa \Rightarrow ab = ba$. Hence S is commutative, and since every commutative AG-groupoid is associative. Therefore, S is commutative semigroup. Finally,

$(vi) \Rightarrow (i)$. Obvious.

Hence the theorem is proved. ∎

References

[1] M. A. Kazim and M. Naseerudin, On almost semigroups, *Portugaliae Mathematica,* 36(1) (1977)

[2] M. Polnijo, Transitive groupoids, *Portugaliae Mathematica*, Vol.50, Fase. 1, 62-74 (1993)

[3] V. Peter and N. Stevanovic, AG-test and some general properties of AG-groupoid, *PU.M.A.* volume 6(4) 371-383 (1995)

[4] M. Shah, A theoretical and computational investigation of AG-groups, PhD thesis, Quaid-i-Azam University Islamabad, (2012)

[5] GAP: Groups Algorithm and Programming, Version 4.4.12, 2008, (2012)

[6] A. Distler, M. Shah and V. Sorge, Enumeration of AG-groupoids, *Lecture Notes in Computer Science*, Volume 6824/2011, 1-14 (2011)
[7] Q. Mushtaq and S. M. Yousuf, On Locally Associative LA-semigroup, *J. Nat. Sci. Math.*, XIX(1), 57–62 (1979)
[8] I. Ahmad, M. Rashad and M. Shah, Some New Results on T^1, T^2 and T^4-AG-groupoids, *Res. J. Recent Sciences*, Vol. 2(3), 64-66, March (2013)
[9] J. R. Cho, Pusan, J. Jezek and T. Kepka, Paramedial Groupoids, *Czechoslovak Mathematical Journal*, 49 (124), 277-290 (1996)
[10] M. Shah, I. Ahmad and A. Ali, Discovery of new classes of AG-groupoids, *Res. J. Recent Sciences*, Vol, 1(11), 47-49 (2012)
[11] M. Shah, A. Ali, and I. Ahmad, On introduction of new classes of AG-groupoids, *Res. J. Recent Sciences*, Vol. 2(1), 67-70, January (2013)
[12] M. Rashad, I. Ahmad, Amanullah and M. Shah, On relations between right alternative and nuclear square AG-groupoids, *Int. Math. Forum,* Vol. 8, no. 5 237-243 (2013)
[13] J. Denes, A. D. Keedwell, Latin Squares and their Applications, Academic Press New York and London (1974)

E-mail address: rashad@uom.edu.pk

E-mail address: iahmaad@hotmail.com

Department of Mathematics, University of Malakand, Chakdara Dir(L), Pakistan.

E-mail address: shahmaths_problem@hotmail.com

Department of Mathematics, Government Post Graduate Collage Mardan, Pakistan.